



Diffusive isotope fractionation of deuterated benzene and toluene in aqueous systems

Jin, Biao; Rolle, Massimo

Publication date:
2018

Document Version
Peer reviewed version

[Link back to DTU Orbit](#)

Citation (APA):

Jin, B., & Rolle, M. (2018). *Diffusive isotope fractionation of deuterated benzene and toluene in aqueous systems*. Abstract from CMWR 2018 conference : Computational Methods in Water Resources, Saint-Malo, France.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Diffusive Isotope Fractionation of Deuterated Benzene and Toluene in Aqueous Systems

Biao Jin, State Key Laboratory of Organic Geochemistry, Guangzhou Institute of Geochemistry, Chinese Academy of Sciences, 510640 Guangzhou, China,

Massimo Rolle, Department of Environmental Engineering, Technical University of Denmark, Lyngby 2800, Denmark

Key words: groundwater, organic contaminants, diffusion and isotope fractionation

Introduction

BTEX compounds (i.e., benzene, toluene, ethylbenzene and xylene) are widespread groundwater organic contaminants [1]. Despite the key controlling role of aqueous diffusion for groundwater contaminants transport [2,3], only a few studies have explored the extent and impact of diffusive deuterium isotope fractionation of organic compounds in aqueous solution [4,5]. In this work we experimentally investigate diffusive and dispersive displacement and fractionation of nondeuterated and perdeuterated benzene and toluene [6].

Experimental Setups

Experiments were performed in two distinct setups (Fig. 1), focusing on (i) pure diffusion of the organic compounds under no-flow conditions, and (ii) hydrodynamic dispersion of the different contaminants under flow-through conditions.

1-D Diffusion Experiments. Gel dissection experiments were performed in cylindrical glass tubes using agarose gel (1% w/w). The tubes were sampled by cutting the gel into 1 cm slices with a scalpel. Each slice was sealed in glass vial with screw caps for GC-MS analyses.

2-D Flow-Through Experiments. The flow-through experiments were conducted in a quasi-2-D flow-through chamber filled with homogeneous quartz sand. High-resolution sampling was performed via 10 equally spaced ports at the outlets (1 cm spacing).




Figure 1. Sketch of the setups for the 1-D tube experiments and the 2-D flow-through experiments.

Results

A normal diffusive isotope effect ($D_{C_7D_8}/D_{C_7H_8} = 0.96$) was observed for toluene, resulting in enrichment of the nondeuterated isotopologue in the direction of the diffusive and transverse dispersive fluxes. Conversely, the measured trends for benzene indicated inverse diffusive fractionation ($D_{C_6D_6}/D_{C_6H_6} = 1.02$). Remarkably faster diffusive/dispersive rates were observed for the perdeuterated isotopologue. These outcomes were consistently observed both for the diffusive fronts in the 1-D tubes and at the fringes of the contaminant plumes in the flow-through setup. The unexpected results of this study suggest that diffusive isotope fractionation of organic contaminants cannot be exclusively ascribed to mass and molar volume differences, but the interactions of the compounds with the surrounding water molecules, as well as the structures and patterns of solute hydration play a key role.

References

- [1] T. H. Wiedemeier, H. S. Rifai, C. H. Newell, T. H. Wilson. Natural attenuation of fuels and chlorinated solvents in the subsurface. John Wiley, New York, 617 p. (1999).
- [2] M. Rolle, G. Chiogna, D.L. Hochstetler, P.K. Kitanidis. On the importance of diffusion and compound-specific mixing for groundwater transport: An investigation from pore to field scale, *J. Contam. Hydrol.*, **153**, 51–68, (2013).
- [3] M. Rolle, G. Chiogna, R. Bauer, C. Griebler, P. Grathwohl. Isotopic Fractionation by Transverse Dispersion: Flow-through Microcosms and Reactive Transport Modeling Study. *Environ. Sci. Technol.*, **44**(16), 6167-6173, (2010).
- [4] E.M. LaBolle, G.E. Fogg, J.B. Eweis, J. Gravner, D.G. Leaist. Isotopic fractionation by diffusion in groundwater. *Water Resour. Res.*, **44** (7), (2008).
- [5] B. Jin, M. Rolle, T. Li, S.B. Haderlein. Diffusive fractionation of BTEX and chlorinated ethenes in aqueous solution: quantification of spatial isotope gradients. *Environ. Sci. Technol.* **48**, 6141-6150, (2014).
- [6] B. Jin and M. Rolle, Normal and Inverse diffusive isotope fractionation of deuterated toluene and benzene in aqueous systems. *Environ. Sci. Technol. Letters.*, **4**, 298-304, (2017).